

Data de implementação e rodadas adicionais
2004, 2006, 2013

Cobertura da embalagem

As advertências de saúde precisam cobrir 50% da frente e 50% do verso das embalagens. No geral, 50% do espaço da embalagem é reservado para advertências de saúde.

Freqüência de rotação e histórico

As advertências devem aparecer em um número igual de embalagens durante um período de 24 meses.


Cingapura produziu três rodadas de advertências de saúde. O primeiro conjunto com seis advertências de saúde foi usado de 2004 a 2005. Um segundo conjunto com seis advertências de saúde foi implementado em 2006. Um terceiro conjunto de seis advertências de saúde deve ser implementado 2013.

Restrições sobre informações enganosas

Não especificado.

ADVERTÊNCIAS

2013



ADVERTÊNCIAS

2006



WARNING:
SMOKING INCREASES MISCARRIAGE RISK
QUIT 1800-438-2000



WARNING:
SMOKING CAUSES MOUTH DISEASES
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES 92% OF ORAL CANCERS
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES 92% OF ORAL CANCERS
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES NECK CANCER
QUIT 1800-438-200



WARNING:
SMOKING CAUSES GANGRENE
QUIT: 1800-438-2000

ADVERTÊNCIAS

2004




WARNING:
TOBACCO SMOKE CAN KILL BABIES
QUIT: 1800-438-2000



WARNING:
SMOKING HARMS YOUR FAMILY
QUIT: 1800-438-2000



WARNING:
SMOKING CAN CAUSE A SLOW PAINFUL DEATH
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES LUNG CANCER
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES STROKE
QUIT: 1800-438-2000



WARNING:
SMOKING CAUSES MOUTH DISEASES
QUIT: 1800-438-2000